

BB



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
20.04.2005 Bulletin 2005/16

(51) Int Cl.7: **A61K 38/00, A61K 38/08,
A61K 38/12**

(21) Application number: **00950283.2**

(86) International application number:
PCT/US2000/018217

(22) Date of filing: **29.06.2000**

(87) International publication number:
WO 2001/000224 (04.01.2001 Gazette 2001/01)

(54) **COMPOSITIONS FOR TREATMENT OF SEXUAL DYSFUNCTION**
VERBINDUNGEN ZUR BEHANDLUNG VON SEXUELLEN FUNKTIONSTÖRUNGEN
COMPOSITIONS POUR LE TRAITEMENT DE DYSFONCTIONNEMENTS SEXUELS

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

• **HERBERT, Guy, W.**
Metuchen, NJ (US)

(30) Priority: **29.06.1999 US 142346 P**
05.04.2000 US 194987 P
28.06.2000 US 606501

(74) Representative: **Kupecz, A., Drs. c.s.**
Octrooibureau Los en Stigter B.V.
Weteringschans 96
1017 XS Amsterdam (NL)

(43) Date of publication of application:
17.04.2002 Bulletin 2002/16

(56) References cited:
WO-A-94/22460 WO-A1-99/64002
FR-A- 2 691 465 FR-A- 2 735 131
US-A- 4 649 191 US-A- 5 420 109
US-A- 5 576 290 US-A- 6 051 555

(73) Proprietor: **Palatin Technologies, Inc.**
Princeton, NJ 08540 (US)

(72) Inventors:
• **BLOOD, Christine, H.**
Morristown, NJ (US)
• **SHADIACK, Annette, M.**
Sommerset, NJ (US)
• **BERNSTEIN, Joanna, K.**
North Brunswick, NJ (US)

• **LOW M ET AL: "ROLE OF CHAIN TERMINI IN
SELECTIVE STEROIDOGENIC EFFECT OF
ACTH MSH-4-10 ON ISOLATED
ADRENOCORTICAL CELLS" PEPTIDES
(ELMSFORD), vol. 11, no. 1, 1990, pages 29-32,
XP009008137 ISSN: 0196-9781**

EP 1 196 184 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**BACKGROUND OF THE INVENTION**5 **Field of the Invention (Technical Field):**

[0001] The present invention relates to peptide constructs for the treatment of sexual dysfunction in animals, including both male erectile dysfunction and female sexual dysfunction, including methods and formulations for the use and administration of the same.

10

Background Art:

[0002] Note that the following discussion refers to a number of publications by author(s) and year of publication, and that due to recent publication dates certain publications are not to be considered as prior art vis-a-vis the present invention. Discussion of such publications herein is given for more complete background and is not to be construed as an admission that such publications are prior art for patentability determination purposes.

15

[0003] Sexual dysfunction, including both penile erectile dysfunction or impotence and female sexual dysfunction, are common medical problems. Significant effort has been devoted over the last twenty or more years to develop methods, devices and compounds for treatment of sexual dysfunction. While more effort has been undertaken for treatment of penile erectile dysfunction, female sexual dysfunction is also an area to which significant research and effort has been devoted.

20

[0004] At present, one commonly used orally administered drug for treatment of sexual dysfunction in the male is Viagra®, a brand of sildenafil, which is a phosphodiesterase 5 inhibitor, increasing the persistence of cyclic guanosine monophosphate and thereby enhancing erectile response. There are several other medical treatment alternatives currently available depending on the nature and cause of the impotence problem. Some men have abnormally low levels of the male hormone testosterone, and treatment with testosterone injections or pills may be beneficial. However, comparatively few impotent men have low testosterone levels. For many forms of erectile dysfunction, treatment may be undertaken with drugs injected directly into the penis, including drugs such as papaverin, prostaglandin E₁, phenoxylbenzamine or phentolamine. These all work primarily by dilating the arterial blood vessels and decreasing the venous drainage. Urethral inserts, such as with suppositories containing prostaglandin, may also be employed. In addition, a variety of mechanical aids are employed, including constriction devices and penile implants.

25

30

[0005] A variety of treatments have also been explored for female sexual dysfunction, including use of sildenafil, although the Food and Drug Administration has not specifically approved such use. Testosterone propionate has also been employed to increase or augment female libido.

35

[0006] Melanocortin receptor-specific compounds have been explored for use of treatment of sexual dysfunction. In one report, a cyclic α -melanocyte-stimulating hormone (" α -MSH") analog, called Melanotan-II, was evaluated for erectogenic properties for treatment of men with psychogenic erectile dysfunction. Wessells H. et al., *J Urology* 160:389-393 (1998); see also U.S. Patent No. 5,576,290, issued November 19, 1996 to M.E. Hadley, entitled *Compositions and Methods for the Diagnosis and Treatment of Psychogenic Erectile Dysfunction* and U.S. Patent No. 6,051,555, issued April 18, 2000, also to M.E. Hadley, entitled *Stimulating Sexual Response in Females*. The peptides used in U.S. Patent Nos. 5,576,290 and 6,051,555 are also described in U.S. Patent No. 5,674,639, issued October 7, 1997, to V.J. Hruby, M.E. Hadley and F. Al-Obeidi, entitled *Cyclic Analogs of Alpha-MSH Fragments*, and in U.S. Patent No. 5,714,576, issued February 3, 1998, to V.J. Hruby, M.E. Hadley and F. Al-Obeidi, entitled *Linear Analogs of Alpha-MSH Fragments*. Melanotan-II is a peptide of the following formula:

45



50

Additional related peptides are disclosed in U.S. Patent Nos. 5,576,290, 5,674,839, 5,714,576 and 6,051,555. These peptides are described as being useful for both the diagnosis and treatment of psychogenic sexual dysfunction in males and females. These peptides are related to the structure of melanocortins.

55

[0007] In use of Melanotan-II, significant erectile responses were observed, with 8 of 10 treated men developing clinically apparent erections, and with a mean duration of tip rigidity greater than 80% for 38 minutes with Melanotan-II compared to 3.0 minutes with a placebo ($p = 0.0045$). The drug was administered by subcutaneous abdominal wall injection, at doses ranging from 0.025 to 0.157 mg/kg body weight. Transient side effects were observed, including nausea, stretching and yawning, and decreased appetite.

[0008] The minimum peptide fragment of native α -MSH needed for erectile response is the central tetrapeptide sequence, His⁶-Phe⁷-Arg⁸-Trp⁹ (SEQ ID 1). In general, all melanocortin peptides share the same active core sequence, His-Phe-Arg-Trp, including melanotropin neuropeptides and adrenocorticotropin. Five distinct melanocortin receptor subtypes have been identified, called MC1-R through MC5-R, and of these MC3-R and MC4-R are believed to be expressed in the human brain. MC3-R has the highest expression in the arcuate nucleus of the hypothalamus, while MC4-R is more widely expressed in the thalamus, hypothalamus and hippocampus. A central nervous system mechanism for melanocortins in the induction of penile erection has been suggested by experiments demonstrating penile erection resulting from central intracerebroventricular administration of melanocortins in rats. While the mechanism of His-Phe-Arg-Trp induction of erectile response has not been fully elucidated, it has been hypothesized that it involves the central nervous system, and probably binding to MC3-R and/or MC4-R.

[0009] Other peptides and constructs have been proposed which are ligands that alter or regulate the activity of one or more melanocortin receptors. For example, International Patent Application No. PCT/US99/09216, entitled *Isoquinoline Compound Melanocortin Receptor Ligands and Methods of Using Same*, discloses two compounds that induce penile erections in rats. However, these compounds were administered by injection at doses of 1.8 mg/kg and 3.6 mg/kg, respectively, and at least one compound resulted in observable side effects, including yawning and stretching. Other melanocortin receptor-specific compounds with claimed application for treatment of sexual dysfunction are disclosed in International Patent Application No. PCT/US99/13252, entitled *Spiropiperidine Derivatives as Melanocortin Receptor Agonists*.

[0010] Both cyclic and linear α -MSH peptides have been studied; however, the peptides heretofore evaluated have had an NH₂ group at the carboxyl terminus. See, for example, Wessells H. et al., *J Urology*, cited above; Haskell-Luevano C. et al., *J Med Chem* 40:2133-39 (1997); Schiöth H.B. et al., *Brit J Pharmacol* 124:75-82 (1998); Schiöth H. B. et al., *Eur J Pharmacol* 349:359-66 (1998); Hadley M.E. et al., *Pigment Cell Res* 9:213-34 (1996); Bednarek M.A. et al., *Peptides* 20:401-09 (1999); U.S. Patent Nos. 6,054,556, 6,051,555 and 5,576,290; and, International Patent Applications PCT/US99/04111 and PCT/US98/03298. While significant research has been conducted in an effort to determine the optimal structure of α -MSH peptides, including a variety of structure-function, agonist-antagonist, molecular modeling and pharmacophore studies, such studies have relied upon peptides with an art conventional NH₂ group at the carboxyl terminus. Further, it has long been believed that biologically active neuropeptides, including α -MSH peptides, are amidated, with an NH₂ group at the carboxyl terminus, and that such amidation is required both for biological activity and stability. See, for example, *Metabolism of Brain Peptides*, Ed. G. O'Cuinn, CRC Press, New York, 1995, pp. 1-9 and 99-101.

[0011] The reference FR-A-2 735 131 discloses conjugates of α MSH with a fatty acid, method to prepare them and their use as a drug.

[0012] Furthermore, FR-A-2 691 465 relates to complexes comprising at least a peptide derived from α MSH, peptide, microsphere, medication and galenic composition comprising the same.

SUMMARY OF THE INVENTION (DISCLOSURE OF THE INVENTION)

[0013] The invention relates to a peptide that is a free acid or pharmaceutically acceptable salt thereof that is Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.

[0014] The invention also includes pharmaceutical compositions of matter, including a peptide of this invention and a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier may be a buffered aqueous carrier, and preferably a saline or citrate buffered carrier.

[0015] The peptide of this invention, and pharmaceutical compositions of this invention, may be for treatment of sexual dysfunction in a mammal. The composition comprises a peptide or pharmaceutically acceptable salt thereof of the formula Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH. The mammal may be male or female. The composition can also include a pharmaceutically acceptable carrier. The peptide or pharmaceutical composition may be for administration by any means known in the art, including administration by injection, administration through mucous membranes, buccal administration, oral administration, dermal administration, inhalation administration and nasal administration. In a preferred embodiment, administration is by nasal administration of a metered amount of a formulation including an aqueous buffer, which buffer may be a saline or citrate buffer.

[0016] A primary object of the present invention is a melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction.

[0017] A second object is to provide a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of male sexual dysfunction, including erectile dysfunction.

[0018] Yet another object is to provide a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of female sexual dysfunction.

[0019] Yet another object is to provide a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction which is effective by intravenous administration at doses of less than 25 μ g/kg of body

weight, and preferably at doses of less than 0.5 µg/kg of body weight.

[0020] Yet another object is to provide a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction which is effective by nasal administration at doses of less than 75 µg/kg of body weight, and preferably at doses of less than 25 µg/kg of body weight.

5 [0021] Yet another object is to provide a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction with substantially reduced incidence of undesirable side effects.

[0022] A primary advantage of the present invention is that it is therapeutically effective at doses that do not cause deleterious side effects, such side effects including nausea, yawning, stretching, decreased appetite and other effects observed with Melanotan-II.

10 [0023] A second advantage of the present invention is that it provides compositions with a larger therapeutic window between desired therapeutic effects and the onset of undesired side effects than other melanocortin receptor-specific agents for the intended purpose.

[0024] Yet another advantage of the present invention is that it provides compositions with a greater safety margin between desired therapeutic effects and the onset of undesired side effects than other melanocortin receptor-specific agents for the intended purpose.

15 [0025] Yet another advantage of the present invention is that it provides a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction which is efficacious at significantly lower doses than Melanotan-II or other melanocortin receptor-specific agents.

[0026] Yet another advantage of the present invention is that it provides a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction which is pharmaceutically active more rapidly following administration than Melanotan-II or other peptide-based melanocortin receptor-specific agents.

20 [0027] Yet another advantage of the present invention is that it provides a peptide-based melanocortin receptor-specific pharmaceutical for use in treatment of sexual dysfunction which, because of increased efficacy at low doses, may be administered by delivery systems other than art conventional intravenous, subcutaneous or intramuscular injection, including but not limited to nasal delivery systems and mucous membrane delivery systems.

25 [0028] Other objects, advantages and novel features, and further scope of applicability of the present invention will be set forth in part in the detailed description to follow, taken in conjunction with the accompanying drawings. The objects and advantages of the invention may be realized and attained by means of the instrumentalities and combinations particularly pointed out in the appended claims.

30

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] The accompanying drawings, which are incorporated into and form a part of the specification, illustrate several embodiments of the present invention and, together with the description, serve to explain the principles of the invention.

35 The drawings are only for the purpose of illustrating a preferred embodiment of the invention and are not to be construed as limiting the invention. In the drawings:

FIG. 1 is a plot of the pharmacokinetic profile in rats intravenously administered 100 µg/kg body weight of **Compound 1**;

40

FIG. 2 is a plot of the pharmacokinetic profile in monkeys intravenously administered 50 µg/kg body weight of **Compound 1**;

FIG. 3 is a plot of the pharmacokinetic profile in a monkey intranasally administered 50 µg/kg body weight of **Compound 1**;

45

FIG. 4 is a plot of the intranasal efficacy in rats of different amounts of **Compound 1**; and

FIG. 5 is a plot of pharmacokinetic profile in Beagle dogs intranasally administered 30, 150 and 650 µg/kg body weight of **Compound 1**.

50

DESCRIPTION OF THE PREFERRED EMBODIMENTS

(BEST MODES FOR CARRYING OUT THE INVENTION)

55

[0030] In the listing of compounds according to the present invention, the amino acid residues have their conventional meaning as given in Chapter 2400 of the Manual of Patent Examining Procedure, 7th Ed. Thus, "Nle" is norleucine; "Asp" is aspartic acid; "His" is histidine; "D-Phe" is D-phenylalanine; "Arg" is arginine; "Trp" is tryptophan; "Lys" is lysine;

"Gly" is glycine; "Pro" is proline; "Tyr" is tyrosine, and "Ser" is serine. "Ac" refers to a peptide or amino acid sequence that is acetylated.

[0031] The invention provides a deamidate α -MSH peptide, that includes the core α -MSH sequence His-D-Phe-Arg-Trp, which is to say that it does not include an $-NH_2$ group at the carboxyl terminus, and has an $-OH$ group at the carboxyl terminus.

[0032] Thus, the invention provides the peptide



Compound 1

[0033] The peptide of **Compound 1** has a formula of $C_{50}H_{68}N_{14}O_{10}$, and a net molecular weight of 1025.18. This peptide may be synthesized by solid-phase means and purified to greater than 96% purity by HPLC, yielding a white powder that is a clear, colorless solution in water.

[0034] In general, the peptide compounds of this invention may be synthesized by solid-phase synthesis and purified according to methods known in the art. Any of a number of well-known procedures utilizing a variety of resins and reagents may be used to prepare the compounds of this invention.

[0035] The peptides of this invention may be in the form of any pharmaceutically acceptable salt. Acid addition salts of the compounds of this invention are prepared in a suitable solvent from the peptide and an excess of an acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, trifluoroacetic, maleic, succinic or methanesulfonic. The acetate salt form is especially useful. Where the compounds of this invention include an acidic moiety, suitable pharmaceutically acceptable salts may include alkali metal salts, such as sodium or potassium salts, or alkaline earth metal salts, such as calcium or magnesium salts.

[0036] The invention provides a pharmaceutical composition that includes a peptide of this invention and a pharmaceutically acceptable carrier. The carrier may be a liquid formulation, and is preferably a buffered, isotonic, aqueous solution. Pharmaceutically acceptable carriers also include excipients, such as diluents, carriers and the like, and additives, such as stabilizing agents, preservatives, solubilizing agents, buffers and the like, as hereafter described.

[0037] **Routes of Administration.** Because, in part, of the increased potency of **Compound 1** of this invention, these peptides may be administered by means other than by injection. If it is administered by injection, the injection may be intravenous, subcutaneous, intramuscular, intraperitoneal or other means known in the art. The peptides of this invention may be formulated by any means known in the art, including but not limited to formulation as tablets, capsules, caplets, suspensions, powders, lyophilized preparations, suppositories, ocular drops, skin patches, oral soluble formulations, sprays, aerosols and the like, and may be mixed and formulated with buffers, binders, excipients, stabilizers, anti-oxidants and other agents known in the art. In general, any route of administration by which the peptides of invention are introduced across an epidermal layer of cells may be employed. Administration means may induce administration through mucous membranes, buccal administration, oral administration, dermal administration, inhalation administration, nasal administration and the like. The dosage for treatment of male erectile dysfunction is administration, by any of the foregoing means or any other means known in the art, of an amount sufficient to bring about an erection of the penis in a male. The dosage for treatment of female sexual dysfunction is administration, by any of the foregoing means or any other means known in the art, of an amount sufficient to bring about the desired response.

[0038] The peptides of this invention may be formulated or compounded into pharmaceutical compositions that include at least one peptide of this invention together with one or more pharmaceutically acceptable carriers, including excipients, such as diluents, carriers and the like, and additives, such as stabilizing agents, preservatives, solubilizing agents, buffers and the like, as may be desired. Formulation excipients may include polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate. For injection or other liquid administration formulations, water containing at least one or more buffering constituents is preferred, and stabilizing agents, preservatives and solubilizing agents may also be employed. For solid administration formulations, any of a variety of thickening, filler, bulking and carrier additives may be employed, such as starches, sugars, fatty acids and the like. For topical administration formulations, any of a variety of creams, ointments, gels, lotions and the like may be employed. For most pharmaceutical formulations, non-active ingredients will constitute the greater part, by weight or volume, of the preparation. For pharmaceutical formulations, it is also contemplated that any of a variety of measured-release, slow-release or time-release formulations and additives may be employed, so that the dosage may be formulated so as to effect delivery of a peptide of this invention over a period of time.

[0039] In general, the actual quantity of peptides of this invention administered to a patient will vary between fairly wide ranges depending upon the mode of administration, the formulation used, and the response desired.

[0040] **Nasal or Intrapulmonary Administration.** By "nasal administration" is meant any form of intranasal administration of any of the peptides of this invention. The peptides may be in an aqueous solution, such as a solution including saline, citrate or other common excipients or preservatives. The peptides may also be in a dry or powder formulation.

[0041] In an alternative embodiment, peptides of this invention may be administered directly into the lung. Intrapulmonary administration may be performed by means of a metered dose inhaler, a device allowing self-administration of a metered bolus of a peptide of this invention when actuated by a patient during inspiration.

5 [0042] The peptides of this invention may be formulated with any of a variety of agents that increase effective nasal absorption of drugs, including peptide drugs. These agents should increase nasal absorption without unacceptable damage to the mucosal membrane. U.S. Patents No. 5,693,608, 5,977,070 and 5,908,825, among others, teach a number of pharmaceutical compositions that may be employed, including absorption enhancers, and the teachings of each of the foregoing, and all references and patents cited therein, are incorporated by reference.

10 [0043] If in an aqueous solution, the peptide may be appropriately buffered by means of saline, acetate, phosphate, citrate, acetate or other buffering agents, which may be at any physiologically acceptable pH, generally from about pH 4 to about pH 7. A combination of buffering agents may also be employed, such as phosphate buffered saline, a saline and acetate buffer, and the like. In the case of saline, a 0.9% saline solution may be employed. In the case of acetate, phosphate, citrate, acetate and the like, a 50 mM solution may be employed. In addition to buffering agents, a suitable preservative may be employed, to prevent or limit bacteria and other microbial growth. One such preservative that may
15 be employed is 0.05% benzalkonium chloride.

[0044] It is also possible and contemplated that the peptide may be in a dried and particulate form. In a preferred embodiment, the particles are between about 0.5 and 6.0 μm , such that the particles have sufficient mass to settle on the lung surface, and not be exhaled, but are small enough that they are not deposited on surfaces of the air passages prior to reaching the lung. Any of a variety of different techniques may be used to make dry powder microparticles, including but not limited to micro-milling, spray drying and a quick freeze aerosol followed by lyophilization. With microparticles, the peptides may be deposited to the deep lung, thereby providing quick and efficient absorption into the bloodstream. Further, with such approach penetration enhancers are not required, as is sometimes the case in transdermal, nasal or oral mucosal delivery routes. Any of a variety of inhalers can be employed, including propellant-based aerosols, nebulizers, single dose dry powder inhalers and multidose dry powder inhalers. Common devices in current
20 use include metered dose inhalers, which are used to deliver medications for the treatment of asthma, chronic obstructive pulmonary disease and the like. Preferred devices include dry powder inhalers, designed to form a cloud or aerosol of fine powder with a particle size that is always less than about 6.0 μm . One type of dry powder inhaler in current use is Glaxo's Rotahaler™, which dispenses a unit dose of powder into a tube, and employs patient suction for inhalation of the powder. Other, more advanced and preferred dry powder inhalers have been or are in development, which
25 include propellants and the like.

[0045] Microparticle size, including mean size distribution, may be controlled by means of the method of making. For micro-milling, the size of the milling head, speed of the rotor, time of processing and the like control the microparticle size. For spray drying, the nozzle size, flow rate, dryer heat and the like control the microparticle size. For making by means of quick freeze aerosol followed by lyophilization, the nozzle size, flow rate, concentration of aerosoled solution and the like control the microparticle size. These parameters and others may be employed to control the microparticle
30 size.

[0046] In one preferred embodiment, a dry powder inhaler is employed which includes a piezoelectric crystal that deaggregates a dry powder dose, creating a small powder "cloud." Once the powder cloud is generated, an electrico-statically charged plate above the powder cloud lifts the drug into the air stream. The user with one relatively easy
40 breath can then inhale the powder. The device may be breath activated, utilizing a flow sensor that activates the electronic components upon the start of inhalation, and thereby eliminating the need for coordination of activation and breathing rhythms by the user.

[0047] Induction of Penile Response. Both Compound 1 and Melanotan-II induce penile erections in experimental rat models when administered by intravenous routes and by other routes, including nasal administration. However,
45 Compound 1 is approximately 100-fold more potent than Melanotan-II for inducing penile erection in the rat model. In intravenous dosing studies, penile erection was induced by Compound 1 in a broad range of concentrations, from 0.5 $\mu\text{g/kg}$ of body weight to 25 $\mu\text{g/kg}$ of body weight. At intravenous doses of 2 $\mu\text{g/kg}$ of body weight, Compound 1 resulted in penile erection in 100% of rats tested, with approximately 50% of the rats having multiple erections during a 30 minutes observation period. By comparison, the optimal efficacious dose intravenously of Melanotan-II in the
50 same rat model was 100 $\mu\text{g/kg}$ of body weight.

[0048] The therapeutic window (the range from the desired therapeutic effect to observed adverse effects) for Compound 1 is on the order of >1,000-fold, compared to a 3- to 4-fold therapeutic window for Melanotan-II. That is, the optimal efficacious dose of Melanotan-II is approximately one-third to one-fourth the dose that causes observed adverse effects. For Compound 1, the optimal efficacious dose is less than one-one thousandth the dose that causes observed
55 adverse effects. The significantly greater therapeutic window for Compound 1 demonstrates that it is better tolerated than Melanotan-II.

[0049] The potency of Compound 1 is significantly higher than that of Melanotan-II, meaning that significantly less Compound 1 is required for a desired effect, as compared to Melanotan-II. For example, 2 $\mu\text{g/kg}$ of body weight of

Compound 1 resulted in penile response by intravenous injection in rats, while 100 µg/kg of body weight of Melanotan-II was required for an equivalent penile response. The significantly higher potency results in less product being required. It further permits utilization of alternative delivery routes, including dermal, nasal and similar delivery routes, wherein higher quantities of drug may be required in order to achieve the desired effect. For example, intranasal routes of administration typically have a bioavailability substantially less than that achieved with intravenous dosing, and thus more drug must be administered by intranasal routes in order to achieve the equivalent pharmacological response.

[0050] Significant commercial advantage for **Compound 1** is demonstrated by the reduced effective dose, compared to Melanotan-II, as a treatment or diagnostic for erectile dysfunction. Among the potential commercial advantages of **Compound 1** are improved effectiveness; lower costs of the product; improved convenience, particularly in that the lower dose permits non-conventional delivery methods that are not feasible with Melanotan-II doses; and improved safety with a decreased risk of side-effects.

Industrial Applicability:

[0051] The invention is further illustrated by the following non-limiting examples.

EXAMPLE 1 Peptide Synthesis

[0052] The peptide Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH was synthesized by standard solid phase peptide synthesis methods, and is a cyclic heptapeptide melanocortin peptide analog with a free acid at the carboxyl terminus and an acetylated amino group at the amino terminus, with the structure:



The peptide has a net molecular weight of 1025.18, and is supplied in an acetate salt form. The peptide is a white, odorless amorphous hygroscopic powder, soluble in 0.9% saline, composed of $\text{C}_{50}\text{H}_{68}\text{N}_{14}\text{O}_{10}$. For synthesis, an Fmoc-Lys(R³)-p-alkoxybenzyl alcohol resin was transferred to a solid phase peptide synthesizer reactor with a mechanical stirrer. The R³ group, such as 1-(1'-adamantyl)-1-methyl-ethoxycarbonyl (Adpoc), allyloxycarbonyl (Aloc) or 4-methyl-trityl (Mtt), was removed and the next Fmoc-protected amino acid (Fmoc-Trp(Boc)-OH) was added to the resin through standard coupling procedures. The Fmoc protective group was removed and the remaining amino acids added individually in the correct sequence, by repeating coupling and deprotection procedures until the amino acid sequence was completed. After completion of coupling with the last Fmoc-amino acid derivative, Fmoc-Nle-OH, and cleavage of the Fmoc protective group, the exposed terminal amino group was acetylated with acetic anhydride and pyridine in N,N-dimethylformamide (DMF). The peptide-resin was dried and the Lys and Asp protective groups cleaved. The Lys and Asp deprotected peptide resin was suspended in a suitable solvent, such as DMF, dichloromethane (DCM) or 1-methyl-2-pyrrolidone (NMP), a suitable cyclic coupling reagent, such as 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TBTU), 2-(7-azo-1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TATU), 2-(2-oxo-1(2H)-pyridyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TPTU) or N,N'-dicyclohexylcarbodiimide/1-hydroxybenzotriazole (DCCI/HOBT) was added, and coupling initiated by use of a suitable base, such as N,N-diisopropylethylamine (DIPEA), sym-collidine or N-methylmorpholine (NMM). After cyclization, the peptide-resin was washed and the peptide cleaved from the resin and any remaining protective groups using trifluoroacetic acid (TFA) in the presence of water and 1,2-ethanedithiol (EDT). The final product was precipitated by adding cold ether and collected by filtration. Final purification was by reversed phase HPLC using a C₁₈ column. The purified peptide was converted to acetate salt by passage through an ion-exchange column.

EXAMPLE 2 Binding of Compound 1 and Melanotan-II to Melanocortin Receptors

[0053] The binding of **Compound 1**, made as in Example 1 above, to selected melanocortin receptors was compared to binding of Melanotan-II. Melanotan-II was also made by conventional peptide synthesis, as generally described in Example 1 above.

[0054] Relative binding was determined by competitive inhibition using α-MSH. B16-F1 melanoma cells were used as the source of MC1 receptors; HEK 293 cells, transfected with human melanocortin receptor sequences, were used as the source of MC3, MC4 and MC5 receptors; and Y1 cells were used as the source of MC2 receptors. A standard competitive binding assay protocol was following, using ¹²⁵I-NDP-MSH as the radioligand.

[0055] Utilizing the assay procedure, the following relative binding was determined:

TABLE 1

RELATIVE BINDING (KI)					
COMPOUND	MC1-R (nM)	MC2-R	MC3-R (nM)	MC4-R (nM)	MC5-R (nM)
Compound 1	0.68	>10 ⁻⁶ M	72.07	19.25	166.8
Melanotan-II	0.12	>10 ⁻⁶ M	23.7	1.03	69.48

EXAMPLE 3 Agonist Activity of Compound 1

[0056] The agonist activity of **Compound 1**, made as in **Example 1** above, was evaluated by cyclic adenosine 3': 5'-cyclic monophosphate (cAMP) detection using a commercially available cAMP kit (R&D Systems, DE0350, low pH). HEK 293 cells, stably transfected with human MC4-R, were used. **Compound 1** was evaluated by addition of each to cells both with and without concomitant addition of α -MSH, with α -MSH also used as a positive control. Following incubation, the medium was aspirated and cell layers extracted. The total cAMP accumulation in fixed quantities of cell extract was determined by competitive immunoassay using the cAMP kit. These results are shown on **Table 2**.

TABLE 2

cAMP Concentration (pMol/ml)				
Concentration of Test Compound	Compound 1	Compound 1 + 100 nM α -MSH	α -MSH	Blank (DMEM)
1 x 10 ⁻⁷ M	270.2	180.9	107.7	8.24
2.5 x 10 ⁻⁸ M	85.1	218.5	146.3	1.9
6.25 x 10 ⁻⁹ M	115.6	62.6	45.0	6.5
1.56 x 10 ⁻⁹ M	17.9	245.8	107.7	0.7
3.91 x 10 ⁻¹⁰ M	22.7	127.0	20.2	2.6
0.98 x 10 ⁻¹⁰ M	5.8	251.8	40.0	0.5

These results demonstrate that **Compound 1** exhibits agonist activity to MC4-R.

EXAMPLE 4 Induction of Penile Erections Using Compound 1 and Melanotan-II

[0057] The efficacious dose of **Compound 1**, made as in **Example 1** above, to induce penile erection in Sprague Dawley rats by intravenous dosing was determined. In the rat model of penile erection, the optimal efficacious dose by intravenous administration is 100 μ g/kg body weight. Melanotan-II was also made by conventional peptide synthesis, as generally described in **Example 1** above. Male Sprague Dawley rats were administered **Compound 1**, in the indicated dose, Melanotan-II, or a saline negative control, and were observed for erections and side effects, including excessive grooming, yawning, vacuous chewing, hypoactivity, and heaving. It is known that Melanotan-II induces these side effects as doses as low as 300 μ g/kg body weight. **Table 3** sets forth the results obtained:

Table 3

Induced Penile Erection In Sprague Dawley Rats by Intravenous Injection			
Drug Administered	Dose (μ g/kg)	% Rats with Penile Erection	Number of Erections/ Rat/ 30 Minutes
Compound 1	0.5	50 %	0.5
Compound 1	0.75	50 %	0.5
Compound 1	1.0	100 %	1.0
Compound 1	2.0	100 %	2.0
Compound 1	5.0	100 %	1.8

Table 3 (continued)

Induced Penile Erection In Sprague Dawley Rats by Intravenous Injection			
Drug Administered	Dose ($\mu\text{g/kg}$)	% Rats with Penile Erection	Number of Erections/ Rat/ 30 Minutes
Compound 1	10.0	100 %	2.5
Compound 1	25.0	50 %	0.5
Melanotan-II	10.0	17 %	0.3
Melanotan-II	100.0	75 %	1.0
Saline (Negative Control)	N/A	0 %	0

Two to four rats were observed in each group, with observation over a one-half hour period following administration. No adverse side effects were observed in any rat administered **Compound 1**. In this and comparable experiments, it was observed that **Compound 1** was efficacious at doses as low as 1 $\mu\text{g/kg}$ of body weight when administered by intravenous injection, with efficacy determined by induction of penile erection in 100% of animals administered **Compound 1**. The comparable efficacious dose of Melanotan-II, with efficacy also determined by induction of penile erection in 100% of animals, was at least approximately 100 $\mu\text{g/kg}$ body weight. This resulted in the unexpected conclusion that **Compound 1** was at least approximately 50-fold more potent for inducing erection in a rat model when administered by intravenous injection than is Melanotan-II. In other studies, no adverse side effects, including excessive grooming, yawning, vacuous chewing, hypoactivity, or heaving, were observed on administration of intravenous doses of **Compound 1** as high as 3,000 $\mu\text{g/kg}$ body weight.

EXAMPLE 5 Pharmacokinetics in Rats Following IV Administration of Compound 1

[0058] A sensitive bioassay, based on high affinity binding to MCRs, was used to monitor plasma levels of **Compound 1** in dosed rats. Rats were dosed intravenously with 100 $\mu\text{g/kg}$ of **Compound 1**. A bi-phasic pharmacokinetic profile resulted, as shown in FIG. 1.

EXAMPLE 6 Pre-Formulation Studies for Nasal Delivery

[0059] A series of pre-formulation studies for nasal delivery were conducted. This study shows the stability of the peptide Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH, at a concentration of .825 mg/mL, over a period of twelve weeks, with storage at 40° C.

EXAMPLE 7 Formulations for Nasal Delivery

[0060] Based on the pre-formulation study of Example 6, two formulations were prepared. For the first, the peptide Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH was dissolved in a 0.9% saline solution, pH approximately 6.0, at a concentration of .825 mg per mL of solution (Nasal Formulation 1). In an alternative formulation, Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH was dissolved in a 50 mM citrate, pH approximately 6.0, also at a concentration of .825 mg per mL of solution (Nasal Formulation 2).

EXAMPLE 8 Inhibition of Compound 1 Using Melanocortin Antagonist

[0061] The effect of a nonselective melanocortin antagonist, SHU9119 (Hruby VJ, Lu D, Sharma SD, et al. *J Med Chem* 38:3454-3461 (1995)) to inhibit induction of penile erection with **Compound 1** was studied. A group of four Sprague Dawley rats were administered 5 $\mu\text{g/kg}$ of SHU9119 by intravenous administration, and five minutes later 25 or 50 $\mu\text{g/kg}$ of **Compound 1** in Nasal Formulation 1 was administered by nasal administration. No erections were observed in the rats over the one-half hour observation period. No erections were observed in rats administered a saline control by nasal administration. In control groups, 100% of the rats administered the same dose of **Compound 1** by the same route of administration, but without pre-administration of SHU9119, had observed erections. FIG. 4 shows the number of penile erections per rat during the initial 30 minute period post-administration; as shown for the 25 $\mu\text{g/kg}$ level, no penile erections were observed with pre-administration of SHU9119.

EXAMPLE 9 Nasal Administration to Cynomolgus Monkeys

[0062] Nasal Formulation 1 of Example 6 was administered to cynomolgus monkeys by nasal spray, at a dose of approximately 50 µg per kg of body weight. Compared to intravenous injection of the same dose of Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH, the mean bioavailability of the peptide using Nasal Formulation 1 was approximately 31.9% ± 2.7. In another series of experiments, the mean bioavailability of the peptide using Nasal Formulation 1 was 27.9% ± 9.7, and the mean bioavailability of the peptide using Nasal Formulation 2 was 34.7% ± 15.6, in both cases again compared to bioavailability of the same quantity of peptide administered by intravenous injection. In a related cardiovascular safety study using cynomolgus monkeys with intranasal administration of Nasal Formulation 1, an erection lasting approximately one hour was observed in one of four monkeys.

EXAMPLE 10 Pharmacokinetic Profiles Following Administration to Cynomolgus Monkeys

[0063] A sensitive bioassay, based on high affinity binding to MCRs, was used to monitor plasma levels of Compound 1 in cynomolgus monkeys. Monkeys were dosed intravenously with 50 µg/kg of Compound 1. A bi-phasic pharmacokinetic profile resulted, as shown in FIG. 2. A selected monkey was intranasally administered 50 µg/kg of Compound 1 in Nasal Formulation 1, and the resulting pharmacokinetic profile is shown in FIG. 3.

EXAMPLE 11 Nasal Administration Dosing Study in Rats

[0064] In another series of experiments, a dosing study was conducted in rats by nasal administration of from 15 µg per kg of body weight to 1500 µg per kg of body weight, using Nasal Formulation 1. Based on penile erectile response behavior, the peptide was efficacious at all dose amounts administered, and particularly from 25 µg/kg to 75 µg/kg, with pharmacologic effect observed over the range from 15 µg/kg to 1500 µg/kg. In addition, the response rate and side effects differed significantly from those demonstrated with nasal administration of Melanotan-II. With nasal administration of the peptide of Melanotan-II, the penile erection response was primarily in the 2nd and 3rd ten-minute period, while with Nasal Formulation 1 the penile erection response was primarily in the 1st and 2nd ten-minute period. Additionally, no adverse side effects were detected at any dosage level with Nasal Formulation 1, while adverse side effects were seen with administration of higher doses of the peptide of Melanotan-II. Table 4 sets forth the comparative data:

Table 4

Induced Penile Erection in Sprague Dawley Rats by Intranasal Administration ^u			
Drug Administered	Dose (µg/kg)	% Rats with Penile Erection	Number of Erections/ Rat/30 Minutes
Compound 1	15	50%	0.5
Compound 1	25	100 %	2.0
Compound 1	37	100%	2.0
Compound 1	50	100%	2.0
Compound 1	75	100 %	2.5
Compound 1	500	100 %	1.5
Compound 1	1000	100 %	1.5
Compound 1	1500	50 %	0.5
Melanotan-II	100	25 %	0.25
Melanotan-II	500	100 %	1.0
Saline (Negative Control)	N/A	0 %	0

EXAMPLE 12 Pharmacokinetic Profiles Following Administration to Beagle Dogs

[0065] A sensitive bioassay, based on high affinity binding to MCRs, was used to monitor plasma levels of Compound 1 in Beagle dogs. The dogs were administered 0 (control), 30, 150 or 540 µg/kg of Compound 1 in Nasal Formulation 1 by nasal administration. The pharmacokinetic profile through four hours post-administration is shown in FIG. 5.

EXAMPLE 13 Administration For Treatment of Male Erectile Dysfunction

[0066] For treatment of male erectile dysfunction in human males with psychogenic erectile dysfunction, a total dose of 500 µg of **Compound 1** in **Nasal Formulation 1** is administered by means of a nasal administration metered dose device. The device dispenses a 100 µL spray volume, and is administered by the patient into a single nostril. If desired, erectile response is measured using an appropriate device, such as a Rigiscan (Dacomed).

EXAMPLE 14 Administration For Treatment of Female Sexual Dysfunction

[0067] For treatment of female sexual dysfunction in human females with psychogenic sexual dysfunction, a total dose of 500 µg of **Compound 1** in **Nasal Formulation 1** is administered by means of a nasal administration metered dose device. The device dispenses a 100 µL spray volume, and is administered by the patient into a single nostril.
 [0068] The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

SEQUENCE LISTING

[0069]

<110> Palatin Technologies, Inc.

Blood, Christine
 Shadiack, Annette
 Bernstein, Joanna K.
 Herbert, Guy W.

<120> Compositions and Methods for Treatment of Sexual Dysfunction

<130> 70025-PCT-4

<140>

<141>

<150> 60/142,346

<151> 1999-06-29

<150> 60/194,987

< 151 > 2000-04-05

<150> PENDING

<151> 2000-06-28

<160> 1

<170> Patentin Ver. 2.1

<210> 1

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: alpha-melanocyte-stimulation-hormone-tetrapeptide core sequence

<400> 1

**His Phe Arg Trp
1**

5

Claims

1. A peptide consisting of the sequence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 10 2. A composition of matter, comprising a peptide and a pharmaceutically acceptable carrier, said peptide selected from the group consisting of Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH and pharmaceutically acceptable salts of Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 15 3. The composition of matter of claim 2, wherein the peptide consists of the sequence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
4. The composition of matter of claim 2, wherein the pharmaceutically acceptable carrier is a buffered aqueous carrier.
- 20 5. A pharmaceutical composition for treatment of sexual dysfunction in a mammal, comprising a peptide and a pharmaceutically acceptable carrier, said peptide selected from the group consisting of Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH and pharmaceutically acceptable salts of Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 25 6. The pharmaceutical composition of matter of claim 5, wherein the peptide consists of the sequence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
7. The use of a peptide or pharmaceutically acceptable salt thereof of the formula Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH, for the manufacture of a pharmaceutical composition for treatment of sexual dysfunction in a mammal.
- 30 8. The use according to claim 7, wherein the peptide consists of the sequence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
9. The use according to claim 7 or 8, wherein the pharmaceutical composition is a pharmaceutical composition for administration by injection, through mucous membranes, buccally, orally, dermally, by inhalation or by nasal administration.
- 35 10. The use according to claim 9, wherein the pharmaceutical composition is an aqueous buffer-comprising pharmaceutical composition for metered dose administration by nasal administration.
- 40 11. The use according to claim 10, wherein the aqueous buffer is a member selected from the group consisting of saline and citrate buffer.

Patentansprüche

- 45 1. Peptid, das aus der Sequenz Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH besteht.
2. Stoffzusammensetzung, die ein Peptid und einen pharmazeutisch verträglichen Träger umfasst, wobei das Peptid aus der Gruppe ausgewählt ist, die aus Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH und pharmazeutisch verträglichen Salzen von Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH besteht.
- 50 3. Stoffzusammensetzung nach Anspruch 2, worin das Peptid aus der Sequenz Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH besteht.
4. Stoffzusammensetzung nach Anspruch 2, worin der pharmazeutisch verträgliche Träger ein gepufferter wässriger Träger ist.
- 55 5. Pharmazeutische Zusammensetzung zur Behandlung einer sexuellen Dysfunktion bei einem Säugetier, welche ein Peptid und einen pharmazeutisch verträglichen Träger umfasst, wobei das Peptid aus der Gruppe ausgewählt

ist, die aus Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH und pharmazeutisch verträglichen Salzen von Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH besteht.

- 5 6. Pharmazeutische Stoffzusammensetzung nach Anspruch 5, worin das Peptid aus der Sequenz Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH besteht.
7. Verwendung eines Peptids mit der Formel Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH oder eines pharmazeutisch verträglichen Salzes davon zur Herstellung einer pharmazeutischen Zusammensetzung für die Behandlung einer sexuellen Dysfunktion bei einem Säugetier.
- 10 8. Verwendung nach Anspruch 7, wobei das Peptid aus der Sequenz Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH besteht.
9. Verwendung nach Anspruch 7 oder 8, wobei die pharmazeutische Zusammensetzung eine für eine Verabreichung ist, die oral, transdermal, durch Mundhöhle, Nase und eine Schleimhaut durch Injektion oder Inhalation erfolgt.
- 15 10. Verwendung nach Anspruch 9, wobei die pharmazeutische Zusammensetzung eine einen wässrigen Puffer enthaltende pharmazeutische Zusammensetzung zur dosierten Verabreichung durch die Nase ist.
- 20 11. Verwendung nach Anspruch 10, wobei der wässrige Puffer aus der Gruppe ausgewählt ist, die aus Citrat- und physiologischem Kochsalzlösungspuffer besteht.

Revendications

- 25 1. Peptide constitué de la séquence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
2. Composition comprenant un peptide et un excipient pharmaceutiquement acceptable, ledit peptide étant choisi dans le groupe se composant de Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH et des sels pharmaceutiquement acceptables de Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 30 3. Composition selon la revendication 2, dans laquelle le peptide se compose de la séquence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 35 4. Composition selon la revendication 2, dans laquelle l'excipient pharmaceutiquement acceptable est un excipient aqueux tamponné.
5. Composition pharmaceutique pour le traitement de dysfonctionnements sexuels chez un mammifère, comprenant un peptide et un excipient pharmaceutiquement acceptable, ledit peptide étant choisi dans le groupe se composant de Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH et des sels pharmaceutiquement acceptables de Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 40 6. Composition pharmaceutique selon la revendication 5, dans laquelle le peptide se compose de la séquence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
- 45 7. Utilisation d'un peptide ou d'un sel pharmaceutiquement acceptable de celui-ci, de formule Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH, pour la fabrication d'une composition pharmaceutique destinée au traitement de dysfonctionnements sexuels chez un mammifère.
- 50 8. Utilisation selon la revendication 7, dans laquelle le peptide est constitué de la séquence Ac-Nle-cyclo(-Asp-His-D-Phe-Arg-Trp-Lys)-OH.
9. Utilisation selon la revendication 7 ou 8, dans laquelle la composition pharmaceutique est une composition pharmaceutique pour une administration par injection, à travers les muqueuses, de manière buccale, orale, dermique, par inhalation ou par voie nasale.
- 55 10. Utilisation selon la revendication 9, dans laquelle la composition pharmaceutique est une composition pharmaceutique contenant un tampon aqueux pour une administration en doses par voie nasale.

EP 1 196 184 B1

- 11.** Utilisation selon la revendication 10, dans laquelle le tampon aqueux est un élément choisi dans le groupe se composant d'une solution salée ou d'un tampon citrate

5

10

15

20

25

30

35

40

45

50

55

PROFILE IN RATS INTRAVENOUSLY ADMINISTERED




FIG. 1

PROFILE IN MONKEYS INTRAVENOUSLY ADMINISTERED






FIG. 2


PROFILE IN MONKEYS INTRANASALLY DELIVERED**FIG. 3**




INTRANASAL EFFICACY OF COMPOUND 1 IN RATS

FIG. 4

Compound 1 Plasma Levels in Beagle Dogs (Intranasal)

**FIG. 5**



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 196 184 B8**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

Note: Bibliography reflects the latest situation

(15) Correction information:
Corrected version no 1 (W1 B1)
INID code(s) 73

(51) Int Cl.7: **A61K 38/00, A61K 38/08,
A61K 38/12**

(48) Corrigendum issued on:
15.06.2005 Bulletin 2005/24

(86) International application number:
PCT/US2000/018217

(45) Date of publication and mention
of the grant of the patent:
20.04.2005 Bulletin 2005/16

(87) International publication number:
WO 2001/000224 (04.01.2001 Gazette 2001/01)

(21) Application number: **00950283.2**

(22) Date of filing: **29.06.2000**

(54) **COMPOSITIONS FOR TREATMENT OF SEXUAL DYSFUNCTION**

VERBINDUNGEN ZUR BEHANDLUNG VON SEXUELLEN FUNKTIONSTÖRUNGEN

COMPOSITIONS POUR LE TRAITEMENT DE DYSFONCTIONNEMENTS SEXUELS

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

- **BERNSTEIN, Joanna, K.**
North Brunswick, NJ (US)
- **HERBERT, Guy, W.**
Metuchen, NJ (US)

(30) Priority: **29.06.1999 US 142346 P**
05.04.2000 US 194987 P
28.06.2000 US 606501

(74) Representative: **Kupecz, A., Drs. c.s.**
Octroolbureau Los en Stigter B.V.
Weteringschans 96
1017 XS Amsterdam (NL)

(43) Date of publication of application:
17.04.2002 Bulletin 2002/16

(60) Divisional application:
05075914.1

(56) References cited:
WO-A-94/22460 WO-A1-99/64002
FR-A- 2 691 465 FR-A- 2 735 131
US-A- 4 649 191 US-A- 5 420 109
US-A- 5 576 290 US-A- 6 051 555

(73) Proprietor: **Palatin Technologies, Inc.**
Cranbury, NJ 08512 (US)

(72) Inventors:
• **BLOOD, Christine, H.**
Morristown, NJ (US)
• **SHADIACK, Annette, M.**
Sommerset, NJ (US)

- **LOW M ET AL: "ROLE OF CHAIN TERMINI IN
SELECTIVE STEROIDOGENIC EFFECT OF
ACTH MSH-4-10 ON ISOLATED
ADRENOCORTICAL CELLS" PEPTIDES
(ELMSFORD), vol. 11, no. 1, 1990, pages 29-32,
XP009008137 ISSN: 0196-9781**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 196 184 B8